

are submerged above 100 C. by increasing the pressure under which ebullition is effected, beyond the normal atmospheric limit.

A fourth point to which we gave attention was the possible preservative effect of "lumps" on Bacteria or their germs. No one would have supposed that Dr. Bastian neglected the precaution of removing large particles of cheese from his experimental infusion. We always strained our cheese emulsion very carefully, or else filtered it. Prof. Cohn found that an infusion made by boiling a pea in water developed Bacteria when the pea was left in it; but if the pea were removed, and the infusion subsequently reboiled, no Bacteria were developed. We found that lumps of cheese could really act as protective hiding-places for Bacterian contamination. In a retort—similar in every respect to Dr. Bastian's—this result was first obtained, though other retorts similarly treated were barren. Accordingly we prepared twelve tubes exactly alike, with the exception that in six the cheese was added as an emulsion, in the other six in the form of lumps. The tubes were closed, and submerged in boiling water for five minutes. Of the "emulsion"-tubes, one burst in the boiling, the other five were barren; of the "lumpy"-tubes, four developed Bacteria in quantity, two remained barren.

In the experiments recorded in NATURE, vol. viii. p. 141, by Dr. Sanderson, it is shown that even when "lumps" are avoided, and the infusion heated by submergence in boiling water, this may not prevent the development of Bacteria when a large bulk of material is employed. But boiling for such a length of time as one hour, or heating to 101° C., always gave him a barren infusion. Dr. Sanderson does not believe that there is a definite relation between the precise temperature to which the infusion is exposed and the destruction of Bacterian contamination, but that the longer heating, or the heating to a higher degree, will increase the chance that Bacteria or their germs are destroyed. Further, Dr. Sanderson's results agree with those of Dr. Pöde and myself as to simple turnip infusion. With this infusion I understand that he has not found the same length or amount of heating necessary as with the turnip infusion to which a fragment of cheese has been added.

And now, I wish very briefly to point out where Dr. Bastian's statements are affected by these results. It is necessary that this should be clearly and simply put, because I find that many persons are under the impression that the investigation of the grounds of Dr. Bastian's statements has shown that there was some solid foundation for them. This is, however, in my opinion, not the case. It is *not* "beyond all question of doubt or cavil that living Bacteria, Torulæ, and other low forms of life will make their appearance and multiply within hermetically-sealed flasks (containing organic infusions) which had been previously heated to 212° F. even for one or two hours." On the contrary, no organic nor inorganic infusion has been contrived by Dr. Bastian nor by anyone else which will develop Bacteria, still less Torulæ, after exposure for one hour (or even less) to 212° F. This is the conclusion given by the impartial examination of the subject, indicated in the experiments above quoted.

Moreover, the statement in the second quotation from Dr. Bastian is abundantly contradicted by the experience of Dr. Sanderson, Dr. Pöde, and myself. Such a turnip-infusion, placed as directed by Dr. Bastian, does *not* invariably become turbid in one or two days, owing to the presence of myriads of Bacteria. We have often kept such infusions free from Bacteria for many days, and I preserved one in a retort with its beak inclined downwards for more than six months, clear as crystal, but amply capable of sustaining the life of Bacteria, as was proved by its accidental contamination a week ago.

It is my opinion that the only *positive* addition to knowledge which this inquiry about the development of Bacteria in infusions has led to is, that when you have cheese-emulsion, or similar material present in an infusion, you must be a little more careful about heating it than when you have not, if you wish to destroy by the agency of heat the life of Bacteria or their germs contained in the infusion. How it is that cheese-emulsion helps the Bacterian contamination to escape destruction we do not know. Possibly in the same way as the larger lumps do. But that matter remains for inquiry when more is ascertained as to the natural history of the Bacteria. I think we may now feel fully satisfied that "archebiosis" or "abiogenesis" is not in any way rendered more probable than it was before by Dr. Bastian's experiments with organic infusions. Prof. Smith and Mr. Archer, of Dublin—eminent authorities in the study of the lower algæ—have criticised in detail and suggested explanations of some of the statements in the third part of "The Beginnings of Life,"

viz., statements relating to the transformation of various species of organisms into others. They show (the reader may consult Prof. Smith's paper in the October number of the *Quarterly Journal of Microscopical Science*, 1873) that the asserted "facts" of transmutations are *not* facts. It is abundantly demonstrated that the fundamental observations recorded by Dr. Bastian are erroneous, and that he has been mistaken.

Exeter College, Oxford, Sept. 26

E. RAY LANKESTER

Variations of Organs

MY father finds that in his letter, published in your number for September 25, he did not give with sufficient clearness his hypothetical explanation of how useless organs might diminish, and ultimately disappear. I therefore now send you, with his approval, the following further explanation of his meaning.

If one were to draw a vertical line on a wall, and were to measure the heights of several thousand men of the same race against this line, recording the height of each by driving in a pin, the pins would be densely clustered about a certain height, and the density of their distribution would diminish above and below. Quetelet experimentally verified that the density of the pins at any distance above the centre of the cluster was equal to that at a like distance below; he also found that the law of diminution of density on receding from the cluster was given by a certain mathematical expression, to which, however, I need here make no further reference. A similar law obtains, with reference to the circumference of the chest; and one may assume, with some confidence, that under normal conditions, the variation of any organ in the same species may be symmetrically grouped about a centre of greatest density, as above explained.

In what follows I shall, for the sake of brevity, speak of the horns of cattle, but it will be understood that my father considers a like argument as applicable to the variations of any organs of any species in size, weight, colour, capacity for performing a function, &c.

Supposing then that a race of cattle becomes exposed to unfavourable conditions, my father's hypothesis is that, whilst the larger proportion of the cattle have their horns developed in the same degree as though they had enjoyed favourable conditions, the remainder have their horns somewhat stunted. Now, if we had made a record of the length of horn in the same species under favourable conditions, we should, as in the case of the heights of men, have a central cluster, with a symmetrical distribution of the pins above and below the cluster. According to the hypothesis, the effect of the poor conditions may be represented by the removal of a certain proportion of the pins, taken at hazard, to places lower down, whilst the rest remain *in statu quo*. By this process the central cluster will be slightly displaced downwards, since its upper edge will be made slightly less dense, whilst its lower edge will become denser; and further, the density of distribution will diminish more rapidly above than below the new central cluster.

Now, if horns are useful organs, the cattle with shorter horns will be partially weeded out by natural selection, and will leave fewer offspring; and after many generations of the new conditions, the symmetry of distribution of the pins will be restored by the weeding out of some of those below the cluster, the central cluster itself remaining undisturbed.

If, on the other hand, horns are useless organs, the cattle with stunted horns have as good a chance of leaving offspring (who will inherit their peculiarity) as their long-horned brothers. Thus, after many generations under the poor conditions, with continual intercrossing of all the members, the symmetry of distribution will be again restored, but it will have come about through the general removal of *all* the pins downwards, and this will of course have shifted the central cluster.

If, then, the poor conditions produce a *continuous* tendency to a stunting of the nature above described, there will be two operations going on side by side—the one ever destroying the symmetry of distribution, and the other ever restoring it through the shifting of the cluster downwards.

Thus, supposing the hypothesis to be supported by facts (and my father intends to put this to the test of experiment next summer), there is a tendency for useless organs to diminish and finally disappear, besides those arising from disuse and the economy of nutrition.

Down, Beckenham, Oct. 4

GEORGE H. DARWIN